



BOTOX[®]
onabotulinumtoxinA injection

NO GOAL IS TOO SMALL

*A story for people living
with Spasticity*

INDICATION

BOTOX[®] (onabotulinumtoxinA) is a prescription medicine that is injected into muscles and used to treat increased muscle stiffness in people 2 years of age and older with spasticity. BOTOX has not been shown to help people perform task-specific functions with their upper limbs or increase movement in joints that are permanently fixed in position by stiff muscles.

IMPORTANT SAFETY INFORMATION

BOTOX may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are preexisting before injection. Swallowing problems may last several months.
- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and


cause serious symptoms, including loss of strength and all-over muscle weakness; double vision; blurred vision; drooping eyelids; hoarseness or change or loss of voice; trouble saying words clearly; loss of bladder control; trouble breathing; and trouble swallowing.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.

Please click here for [Consumer Brief Summary](https://www.rxabbvie.com/pdf/botox_pi.pdf) or visit https://www.rxabbvie.com/pdf/botox_pi.pdf for Full Prescribing Information.

ABOUT THIS STORY

This story is intended to provide helpful information about BOTOX® as a treatment option for people living with Spasticity. Botulinum toxin has been studied by scientists for more than 100 years. BOTOX® was approved to treat Adult Upper Limb Spasticity in 2010 and Adult Lower Limb Spasticity in 2016.



IMPORTANT SAFETY INFORMATION (continued)

BOTOX may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of receiving BOTOX. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Do not receive BOTOX if you are allergic to any of the ingredients in BOTOX (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Dysport® (abobotulinumtoxinA), Xeomin® (incobotulinumtoxinA), Jeuveau® (prabotulinumtoxinA-xvfs), Daxxify® (daxibotulinumtoxinA-lanm), or Letybo® (letibotulinumtoxinA-wlbg) (this may not be a complete list of all botulinum toxin products); have a skin infection at the planned injection site.

The dose of BOTOX is not the same as, or comparable to, another botulinum toxin product.

INSPIRED BY REAL PATIENT STORIES

Meet Ellen. Ellen is 48 years old, and her stroke occurred shortly after her 42nd birthday. She was diagnosed with Upper Limb and Lower Limb Spasticity and treats her condition with BOTOX®. Prior to receiving BOTOX®, Ellen was able to walk but occasionally used a wheelchair for ease to move around.


Always consult your doctor before making any treatment decisions.

Spasticity can happen for many different reasons, such as Traumatic Brain Injury (TBI), Multiple Sclerosis (MS), Spinal Cord Injury (SCI), and Cerebral Palsy (CP) but often occurs after a stroke. When you have Spasticity, you may have hands or fingers that are stiff and can't be unclenched; a tight, bent elbow; or curled toes that make it hard to walk.



Please see additional Important Safety Information in the following pages and on pages 26 and 27.

BOTOX[®]
onabotulinumtoxinA_{injection}

- 
- 5 Ellen's BOTOX® story
 - 12 Could it work for me?
 - 15 What about needles?
 - 16 What if I have concerns about BOTOX®?
 - 18 Is BOTOX® a one-time injection?
 - 19 How do I track my progress?
 - 20 How much does BOTOX® cost?
 - 23 Ellen's final thoughts

IMPORTANT SAFETY INFORMATION (continued)

Serious and/or immediate allergic reactions have been reported, including itching; rash; red, itchy welts; wheezing; asthma symptoms; dizziness; or feeling faint. Get medical help right away if you experience symptoms; further injection of BOTOX should be discontinued.



Ellen grabbed her cane and headed to the kitchen for coffee. She felt a slight stiffness in her left wrist and left foot—thank goodness it was almost time for her next BOTOX® appointment.

It'd been nearly 6 years since Ellen was diagnosed with Upper and Lower Limb Spasticity, and she had adjusted well.

She thought about life before her stroke. Did she miss it? Sometimes.

But was she proud of how far she'd come? Absolutely.

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all your muscle or nerve conditions, such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects, including difficulty swallowing and difficulty breathing from typical doses of BOTOX.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.

BOTOX
onabotulinumtoxinA injection



Ellen's stroke happened shortly after her 42nd birthday. In the months that followed, she noticed that the muscles in her left hand and left foot became stiff. Her wrist bent, and her thumb dug into her palms. Sometimes, her toes would clench up too. Getting ready in the morning had become more challenging for her.

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor if you have any breathing-related problems. Your doctor may monitor you for breathing problems during your treatment with BOTOX for spasticity. The risk of developing lung disease in patients with reduced lung function is increased in patients receiving BOTOX.

She remembers telling her doctor that it would mean everything if getting ready for the day was less challenging so that she could feel like less of a burden. So, when he suggested BOTOX® as an option to treat her Spasticity, she was excited to give it a try.

Ellen didn't want her Spasticity to define her.



IMPORTANT SAFETY INFORMATION (continued)

Bronchitis and upper respiratory tract infections (common colds) have been reported. Bronchitis was reported more frequently in adults receiving BOTOX for upper limb spasticity. Upper respiratory infections were also reported more frequently in adults with prior breathing-related problems with spasticity. In pediatric patients treated with BOTOX for upper limb spasticity, upper respiratory tract infections were reported more frequently. In pediatric patients treated with BOTOX for lower limb spasticity, upper respiratory tract infections were not reported more frequently than placebo.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.

BOTOX
onabotulinumtoxinA injection

Ellen sat down next to Paul as he poured coffee into her mug. “Excited for lunch with your friends today?” he asked.

Before Ellen responded, she thought about how much more enjoyable morning coffee with Paul was since starting treatment with BOTOX®.



She smiled, looking down at the warm, steaming mug.

“They haven’t seen me since I started treatment with BOTOX®,” she replied.



IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all your medical conditions, including if you have or have had bleeding problems; have plans to have surgery; had surgery on your face; have weakness of forehead muscles, trouble raising your eyebrows, drooping eyelids, and any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX can harm your unborn baby); are breastfeeding or plan to (it is not known if BOTOX passes into breast milk).



Because of her condition, getting ready took her a little longer. Still, Ellen was proud of her progress with getting ready.



IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you received BOTOX in the past.**

Please see additional Important Safety Information in the following pages and on pages 26 and 27.


BOTOX[®]
onabotulinumtoxinA injection



As her taxi driver headed toward the city, Ellen reflected upon the last time she saw her friends. They all were incredibly supportive, but deep down she felt like an outcast. Ellen didn't like the constant doting and overcompensating from her friends; it was awkward and although it made her feel robbed of her dignity, it also motivated her to talk to her doctor about available treatment options.

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor if you received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as Myobloc®, Dysport®, Xeomin®, Jeuveau®, Daxxify®, or Letybo® in the past (this may not be a complete list of all botulinum toxin products; tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.



Twenty minutes later, Ellen arrived at the restaurant. Marcus and Lisa met her with hugs and kisses.

Once inside, the friends went around the table catching up on all the events that were missed: birthdays, graduations, and even a few health scares. After everyone else had shared, they turned to Ellen.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.



“So, Ellen, how’ve you been? What’s new?”
Marcus asked.

“Well, I’ve started treatment with BOTOX®
for my Spasticity,” Ellen quipped.

“BOTOX®?!” her friends bellowed in unison.

“That can be used for Spasticity?”
Lisa asked.

“Ha! How’d I know you were going to be
surprised.” Ellen said, laughing.

Ellen explained that BOTOX®
injections are locally administered
and work differently than oral
medications when treating Spasticity.




IMPORTANT SAFETY INFORMATION (continued)

Other side effects of BOTOX include dry mouth; discomfort or pain at the injection site; tiredness; headache; neck pain; eye problems such as double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes; drooping eyebrows; and upper respiratory tract infection.

For more information refer to the Medication Guide or talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please click here for [Consumer Brief Summary](#) or visit https://www.rxabbvie.com/pdf/botox_pi.pdf for Full Prescribing Information.



“My doctor injects BOTOX® into the specific muscles within my body that have been affected by Spasticity,” Ellen said, pointing to her left wrist and foot.

Ellen continued to explain that **BOTOX® helps reduce the tightness in her muscles**. Since starting treatment, she’d experienced improvements and was happy she found a way to enjoy one of her favorite hobbies.

“Ah Ell, this is great to hear. Especially since we knew how frustrated you were at times,” Marcus said.

IMPORTANT SAFETY INFORMATION (continued)

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/PatientAccessSupport to learn more.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.

BOTOX®
onabotulinumtoxinA injection



IMPORTANT SAFETY INFORMATION

BOTOX may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are preexisting before injection. Swallowing problems may last several months.



As they polished off their appetizers, Lisa cringed, “I don’t know how you do needles. Don’t they bother you?” Ellen smiled, shaking her head no.

She told them that fine needles are used, but they may cause some discomfort. **“BOTOX® is only injected into my affected muscles.** If any pain occurs where the needle is inserted, it’s temporary,” she added.

Ellen took a sip of water, then finished, “My doctor does it right in his office.”

IMPORTANT SAFETY INFORMATION (continued)

- **Spread of toxin effects.** The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms, including loss of strength and all-over muscle weakness; double vision; blurred vision; drooping eyelids; hoarseness or change or loss of voice; trouble saying words clearly; loss of bladder control; trouble breathing; and trouble swallowing.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.

BOTOX®
onabotulinumtoxinA_{injection}



“Honestly,” she began, “I was hesitant at first because BOTOX® is a toxin. My doctor explained that **BOTOX® is FDA approved to treat both Upper and Lower Limb Spasticity in people 2 years and older.** He also explained what the most common side effects were for Adult Spasticity, which I appreciated, as that’s important for me to know.”

Ellen paused to take a bite of her pasta, then said, “Knowing it’s FDA approved for Spasticity made me feel more comfortable. And you know, not letting my Spasticity define me has been super important to me.”



IMPORTANT SAFETY INFORMATION (continued)

BOTOX may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of receiving BOTOX. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Do not receive BOTOX if you are allergic to any of the ingredients in BOTOX (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Dysport® (abobotulinumtoxinA), Xeomin® (incobotulinumtoxinA), Jeuveau® (prabotulinumtoxinA-xvfs), Daxxify® (daxibotulinumtoxinA-lanm), or Letybo® (letibotulinumtoxinA-wlbg) (this may not be a complete list of all botulinum toxin products); have a skin infection at the planned injection site.

The dose of BOTOX is not the same as, or comparable to, another botulinum toxin product.

MOST COMMON SIDE EFFECTS*

ADULT UPPER LIMB

- NAUSEA
- FATIGUE
- BRONCHITIS
- PAIN IN EXTREMITY
- MUSCULAR WEAKNESS

ADULT LOWER LIMB

- JOINT PAIN
- BACK PAIN
- MUSCLE PAIN
- RESPIRATORY INFECTION
- INJECTION SITE PAIN

“I remember telling my doctor that if I could just get ready with less muscle stiffness, it’d mean a lot to me,” she said.

“So, I gave BOTOX® a try. And let me tell you, I’m so glad I did.”

*These do not cover all the possible serious side effects of BOTOX®. Please see the Consumer Brief Summary about BOTOX® and the Important Safety Information, and talk to your doctor to learn more.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.


BOTOX®
onabotulinumtoxinA injection



Her friends listened intently as she explained that BOTOX® isn't just a one-time treatment. "As my doctor explained," Ellen said, "I go for **BOTOX® injections every 12 weeks** or so. We discuss my progress during each visit and how I'm responding to treatment. My doctor determines how much BOTOX® I need and what muscles need to be injected during those visits, based on how I am doing."

"Believe it or not, I can start to feel when it's time for another injection," Ellen finished.

*BOTOX® treatments may be repeated no sooner than every 12 weeks, once the previous treatment wears off.



IMPORTANT SAFETY INFORMATION (continued)

Serious and/or immediate allergic reactions have been reported, including itching; rash; red, itchy welts; wheezing; asthma symptoms; dizziness; or feeling faint. Get medical help right away if you experience symptoms; further injection of BOTOX should be discontinued.

Tell your doctor about all your muscle or nerve conditions, such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects, including difficulty swallowing and difficulty breathing from typical doses of BOTOX.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.


“You know me,” said Ellen, “I’ve never been the type to keep a journal or a diary.”

“But since starting BOTOX®, **tracking my progress has been super helpful.** I note my improvements, how I’m feeling, questions I have, and any future goals I want to set for myself. It’s kind of awesome, and it really helps me remain focused during each visit with my doctor.”



Learn more about preparing for treatment with BOTOX®.
Visit [BOTOXspasticity.com/what-to-expect](https://www.BOTOXspasticity.com/what-to-expect)

BOTOX[®]
onabotulinumtoxinA injection

An illustration of a restaurant scene. In the foreground, three people are seated at a round white table. A woman with dark hair, wearing a purple top and a light-colored cardigan, is holding up a smartphone and looking at it. Two other people, a man and a woman, are seated across from her, looking towards her. The table has plates of food and glasses. In the background, another table with people is visible, and the floor is light-colored. The overall style is a soft, painterly illustration.

“If you don’t mind me asking, how much does BOTOX® cost?” Lisa asked.

“LISA!!!” Marcus squealed.

“Oh, please. Totally normal question. I mean, we have been friends for 30 years!” Ellen said.

Ellen explained that **cost is different for every individual, depending on eligibility.**

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor if you have any breathing-related problems. Your doctor may monitor you for breathing problems during your treatment with BOTOX for spasticity. The risk of developing lung disease in patients with reduced lung function is increased in patients receiving BOTOX.

Bronchitis and upper respiratory tract infections (common colds) have been reported. Bronchitis was reported more frequently in adults receiving BOTOX for upper limb spasticity. Upper respiratory infections were also reported more frequently in adults with prior breathing-related problems with spasticity. In pediatric patients treated with BOTOX for upper limb spasticity, upper respiratory tract infections were reported more frequently. In pediatric patients treated with BOTOX for lower limb spasticity, upper respiratory tract infections were not reported more frequently than placebo.



“There are options,” Ellen began.

“I have commercial insurance, so most of my out-of-pocket costs are covered. There may be additional costs for the procedure, but really it varies based on healthcare provider and insurance coverage. **There’s actually a pretty cool BOTOX® Savings Program that has helped reimburse some of my remaining costs.**”

“I had no idea. That’s great!” Lisa said with a smile as she flagged the server for the check.

Learn more about ways to save on BOTOX®. Visit [BOTOXspasticity.com/savings](https://www.BOTOXspasticity.com/savings).




You may pay as little as \$0 for out-of-pocket costs not covered by commercial insurance. Text SAVE to 27747* to check your eligibility.

*See Privacy & Terms: [BOTOXSavingsProgram.com/Eligibility](https://www.BOTOXSavingsProgram.com/Eligibility). Message & data rates may apply. Message frequency may vary. Text HELP for help or STOP to end.

Please see additional Important Safety Information in the following pages and on pages 26 and 27.



An illustration showing four people on a sidewalk. On the left, a man in a wheelchair is being embraced by a woman in a white lab coat and a woman in a white top and light pants. To the right, a man in a blue sweater and light pants stands by a dark blue car, looking towards the group. The background features a large green tree and a clear blue sky.

As Ellen and her friends headed toward the exit, they embraced goodbye, promising not to let so much time pass by before doing this again.

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all your medical conditions, including if you have or have had bleeding problems; have plans to have surgery; had surgery on your face; have weakness of forehead muscles, trouble raising your eyebrows, drooping eyelids, and any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX can harm your unborn baby); are breastfeeding or plan to (it is not known if BOTOX passes into breast milk).



On the drive home with her husband, Ellen reflected on the afternoon. “We had a great time. We talked about Lisa’s boys and Marcus’ health scare. But honestly, we talked a lot about my story and the progress I’ve made with BOTOX®.”

She continued, “Maybe one of my friends will pass along this information, and eventually it will make its way to someone who is also living with Spasticity, who can then talk to their doctor to learn more. I forget that not everyone knows that BOTOX® is an option to treat this condition.”

IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you received BOTOX in the past.**

Please see additional Important Safety Information in the following pages and on pages 26 and 27.

BOTOX[®]
onabotulinumtoxinA_{injection}

Ellen looked out the window and smiled. She was proud of her determination and motivation to seek a treatment option that was right for her.

But most of all, she was proud of the fact that she never let Spasticity get in her way.

When living with poststroke Spasticity,

NO GOAL IS TOO SMALL



IMPORTANT SAFETY INFORMATION (continued)

Tell your doctor if you received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as Myobloc®, Dysport®, Xeomin®, Jeuveau®, Daxxify®, or Letybo® in the past (this may not be a complete list of all botulinum toxin products; tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.

Other side effects of BOTOX include dry mouth; discomfort or pain at the injection site; tiredness; headache; neck pain; eye problems such as double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes; drooping eyebrows; and upper respiratory tract infection.



Visit [BOTOXspasticity.com](https://www.botoxspasticity.com) to learn more about treating your Spasticity with BOTOX®



Please see additional Important Safety Information in the following pages and on pages 26 and 27.

BOTOX[®]
onabotulinumtoxinA injection

INDICATION

BOTOX® (onabotulinumtoxinA) is a prescription medicine that is injected into muscles and used to treat increased muscle stiffness in people 2 years of age and older with spasticity. BOTOX has not been shown to help people perform task-specific functions with their upper limbs or increase movement in joints that are permanently fixed in position by stiff muscles.

IMPORTANT SAFETY INFORMATION

BOTOX may cause serious side effects that can be life threatening. Get medical help right away if you have any of these problems any time (hours to weeks) after injection of BOTOX:

- **Problems swallowing, speaking, or breathing**, due to weakening of associated muscles, can be severe and result in loss of life. You are at the highest risk if these problems are preexisting before injection. Swallowing problems may last several months.
- **Spread of toxin effects**. The effect of botulinum toxin may affect areas away from the injection site and cause serious symptoms, including loss of strength and all-over muscle weakness; double vision; blurred vision; drooping eyelids; hoarseness or change or loss of voice; trouble saying words clearly; loss of bladder control; trouble breathing; and trouble swallowing.

BOTOX may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of receiving BOTOX. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Do not receive BOTOX if you are allergic to any of the ingredients in BOTOX (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Dysport® (abobotulinumtoxinA), Xeomin® (incobotulinumtoxinA), Jeuveau® (prabotulinumtoxinA-xvfs), Daxxify® (daxibotulinumtoxinA-lanm), or Letybo® (letibotulinumtoxinA-wlbg) (this may not be a complete list of all botulinum toxin products); have a skin infection at the planned injection site.

The dose of BOTOX is not the same as, or comparable to, another botulinum toxin product.

Serious and/or immediate allergic reactions have been reported, including itching; rash; red, itchy welts; wheezing; asthma symptoms; dizziness; or feeling faint. Get medical help right away if you experience symptoms; further injection of BOTOX should be discontinued.

Tell your doctor about all your muscle or nerve conditions, such as ALS or Lou Gehrig's disease, myasthenia gravis, or Lambert-Eaton syndrome, as you may be at increased risk of serious side effects, including difficulty swallowing and difficulty breathing from typical doses of BOTOX.

Tell your doctor if you have any breathing-related problems. Your doctor may monitor you for breathing problems during your treatment with BOTOX for spasticity. The risk of developing lung disease in patients with reduced lung function is increased in patients receiving BOTOX.

IMPORTANT SAFETY INFORMATION (continued)

Bronchitis and upper respiratory tract infections (common colds) have been reported. Bronchitis was reported more frequently in adults receiving BOTOX for upper limb spasticity. Upper respiratory infections were also reported more frequently in adults with prior breathing-related problems with spasticity. In pediatric patients treated with BOTOX for upper limb spasticity, upper respiratory tract infections were reported more frequently. In pediatric patients treated with BOTOX for lower limb spasticity, upper respiratory tract infections were not reported more frequently than placebo.

Tell your doctor about all your medical conditions, including if you have or have had bleeding problems; have plans to have surgery; had surgery on your face; have weakness of forehead muscles, trouble raising your eyebrows, drooping eyelids, and any other abnormal facial change; are pregnant or plan to become pregnant (it is not known if BOTOX can harm your unborn baby); are breastfeeding or plan to (it is not known if BOTOX passes into breast milk).

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Using BOTOX with certain other medicines may cause serious side effects. **Do not start any new medicines until you have told your doctor that you received BOTOX in the past.**

Tell your doctor if you received any other botulinum toxin product in the last 4 months; have received injections of botulinum toxin such as Myobloc[®], Dysport[®], Xeomin[®], Jeuveau[®], Daxxify[®], or Letybo[®] in the past (this may not be a complete list of all botulinum toxin products; tell your doctor exactly which product you received); have recently received an antibiotic by injection; take muscle relaxants; take an allergy or cold medicine; take a sleep medicine; take aspirin-like products or blood thinners.

Other side effects of BOTOX include dry mouth; discomfort or pain at the injection site; tiredness; headache; neck pain; eye problems such as double vision, blurred vision, decreased eyesight, drooping eyelids, swelling of your eyelids, and dry eyes; drooping eyebrows; and upper respiratory tract infection.

For more information refer to the Medication Guide or talk with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please click here for [Consumer Brief Summary](https://www.rxabbvie.com/pdf/botox_pi.pdf) or visit https://www.rxabbvie.com/pdf/botox_pi.pdf for Full Prescribing Information.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/PatientAccessSupport to learn more.



BOTOX[®]
onabotulinumtoxinA injection



IS SPASTICITY GETTING IN YOUR WAY?

If you're living with Spasticity, treatment should be prioritized. BOTOX® is proven to reduce muscle stiffness.

Talk to your doctor today to see if BOTOX® is the right treatment for you.

IMPORTANT SAFETY INFORMATION (continued)

BOTOX may cause loss of strength or general muscle weakness, vision problems, or dizziness within hours to weeks of receiving BOTOX. **If this happens, do not drive a car, operate machinery, or do other dangerous activities.**

Do not receive BOTOX if you are allergic to any of the ingredients in BOTOX (see Medication Guide for ingredients); had an allergic reaction to any other botulinum toxin product such as Myobloc® (rimabotulinumtoxinB), Dysport® (abobotulinumtoxinA), Xeomin® (incobotulinumtoxinA), Jeuveau® (prabotulinumtoxinA-xvfs), Daxxify® (daxibotulinumtoxinA-lanm), or Letybo® (letibotulinumtoxinA-wlbg) (this may not be a complete list of all botulinum toxin products); have a skin infection at the planned injection site.

Please see additional Important Safety Information, including Boxed Warning, throughout and on pages 26 and 27.

Please click here for [Consumer Brief Summary](https://www.rxabbvie.com/pdf/botox_pi.pdf) or visit https://www.rxabbvie.com/pdf/botox_pi.pdf for Full Prescribing Information.



FIND A BOTOX® SPECIALIST

BOTOXspasticity.com/
#find-a-botox-specialist

SCAN QR CODE



SIGN UP FOR BOTOX® UPDATES

BOTOXspasticity.com/
savings-support

SCAN QR CODE